

97. Some prescription weight-loss drugs have been linked to heart valve problems.
98. Almost one-half of deaths associated with anorexia are from suicide.
99. If you suspect someone of having an eating disorder, you should approach him or her with your concern.
100. There is a vaccine to prevent HIV infection or AIDS.
101. Everyone who is exposed to HIV gets infected.
102. Only men are at risk for HIV infection during vaginal sex.
103. Chlamydia is not curable
104. A person can get HIV from body piercing, ear piercing or tattoo needles.
105. The use of lubricants containing nonoxynol-9 further reduces the risk of HIV transmission.
106. Using a special shampoo can eliminate pubic lice.
107. HIV can be transmitted through human bites.
108. A person should wait for symptoms before getting tested.
109. HIV testing is never mandatory.
110. AIDS is caused by a bacteria.
111. HIV cannot not be transmitted from woman to woman.
112. Wearing tight fitting underwear can cause Herpes outbreaks.
113. All people with AIDS have the same symptoms.
114. Without drugs that help people with AIDS live longer, a person who develops AIDS will usually die within 6 mos. to 2 years.
115. People can become infected with HIV and pass the infection on to a sexual partner without ever having symptoms.
116. Vertical Transmission of HIV is transmission from a man to a woman.
117. Douching after sex can reduce the risk of HIV infection.
118. Sex partners who are both infected with HIV need not use condoms when having sex
119. One is at a higher risk of contracting HIV/AIDS from a 7 year old child than from an infant.
120. A child infected with HIV should receive routine childhood immunizations